



# Porter Medical Associates

## Immunotherapy Dose Schedule

### Dosage and necessary documentation:

Injections are to be recorded by vial, dose, date, and location. Each complete Immunotherapy treatment consists of 5 vials (Phases) and each vial contains 10 doses (10 weeks). The weekly dosage amount is broken down in the diagram below. Injections should be consistent, once a week with a total of 50 weeks at the completion of therapy.

### **Vial#5/Silver, Vial#4/Green, Vial#3/Blue, Vial#2/Gold, Vial#1/Red**

1. First dose .05 cc
2. Second dose .1 cc
3. Third dose .15 cc
4. Fourth dose .2 cc
5. Fifth dose .25 cc
6. Sixth dose .3 cc
7. Seventh dose .35 cc
8. Eighth dose .4 cc
9. Ninth dose .45 cc
10. Tenth dose .5 cc

### What to do if patient misses a dose:

If patient absolutely must miss a dose, they have to go through a build up schedule to

get them back on track and make the therapy effective. **(Missed Doses Not Recommended)**

**-Build-up phase** (time intervals from last injection).

- 0 to 14 days, continue as scheduled
- 15 to 20 days, reduce by 1 dose.
- 21 to 28 days, reduce by 2 doses.
- 29 to 35 days, reduce by 4 doses
- >36 days, contact physician for orders

